

Quantum randomness emerging under gravitational nonlinearity

Tamás Geszti

*Department of the Physics of Complex Systems,
Eötvös University; H-1117 Budapest, Hungary
e-mail: geszti@galahad.elte.hu*

A scenario is outlined for quantum measurement, assuming that self-sustaining classicality is the consequence of an attractive gravitational self-interaction acting on massive bodies, and randomness arises already in the classical domain. A simple solvable model is used to demonstrate that small quantum systems influence big ones in a mean-field way, offering a natural route to Born's probability rule.

PACS numbers: PACS numbers: 03.65.BZ, 42.50.-p

The origin of randomness in quantum mechanics is still unclear, after eighty years since its first recognition by Born[1]. The so-called collapse of the wave function[2], hardly more than a name for the moment when randomness emerges, could not be given a physical identity so far. Von Neumann attached a schematic picture to the name. Assume a measuring apparatus that is capable to distinguish two orthogonal states of a microobject by evolving under interaction with the object into macroscopically distinguishable states. However, if the microobject starts from a superposition of the basis states, a two-act scenario[3] follows: the composite system evolves first unitarily into an entangled superposition of object and apparatus, then non-unitary evolution - identified with wave-function collapse - erases randomly one of the terms of the superposition, resulting in a macroscopically definite outcome. For the simplest case of a two-state microobject with basis states $|+\rangle$ and $|-\rangle$, interacting with a one-dimensional apparatus of center-of-mass (c.o.m.) wave function $\psi(x, t)$, to be analyzed below, the unitary stage of evolution reads

$$\begin{aligned} |\Psi\rangle &= (c_+|+\rangle + c_-|-\rangle)\psi(x, 0) \\ \Rightarrow c_+|+\rangle\psi_+(x, t) + c_-|-\rangle\psi_-(x, t), \end{aligned} \quad (1)$$

$p = |c_+|^2$ and $1-p = |c_-|^2$ being the respective probabilities that in the subsequent non-unitary stage the system would collapse into the first or the second term of the superposition.

It is implicit that the switch from unitary to collapse dynamics takes place somewhere on the borderline between quantum and classical world. Whether the two stages are separated indeed in time, and if so, whether the second can be resolved into more detailed dynamics, remain unanswered. The situation is summarized in the Copenhagen interpretation of quantum mechanics, according to which such questions are declared illegal, the quantum and classical domains are distinct, the former being a mirror world providing dynamics-like rules to predict statistics of random events in the latter[4]. It is one of the surprising facts about Nature that this interpretation works "for all practical purposes", as observed

by John Bell in a tone of dissatisfaction[5].

The theory of environment-induced decoherence[6, 7], although offering an important clue about the quantum-classical border by explaining the gradual disappearance of interference between more and more different states as we traverse the mesoscopic domain, gives no new insight about the nature of collapse. Well tested on its own territory, it seems to offer but a vague clue about collapse: as we learn - in fair agreement between theory and experiment - to keep decoherence at a lower and lower level[8], we seem to observe that in such cases the collapse of the wave function remains also absent: decoherence seems to be a necessary condition for collapse, although the nature of that control - if confirmed - remains to be explained.

A seriously elaborated alternative approach, stochastic reduction models[9] are a fancy phenomenology postulating two new things at a time: a deeper dynamical level with random noise generation, and nonlinear quantum dynamics - no independent evidence seems to be around in favor of the resulting scheme. Below we argue that the theory probably tries to catch too much at one step: randomness need not be put in by hand.

The last remark is based on the observation that for a macroscopic apparatus described by classically chaotic dynamics and coupled to microscopic subsystems in a mean-field fashion instead of quantum mechanics, Born's rule[1] about quantum probabilities would arise in a natural way: chaotic dynamics transform weak forces into probabilities via amplification of small deflections. For short times even linear quantum mechanics acts on wave packets in a mean-field way as exemplified by the theory of so-called weak measurements[10]. Unfortunately, for the long-time phenomenology this fact is of no use: mean-field behavior is intrinsically nonlinear.

Starting with the above motivation, we assume that it suffices to explain the primordial emergence and subsequent persistence of classicality - the resistance of macroscopic bodies to form superpositions; then randomness is generated in the classical domain through some kind of chaotic dynamics. The prospects for such an explanation have been strongly improved through the recently discovered fact that randomness generated in quasi-classical

motion appears in purely linear quantum dynamics as well[11].

The theory of the inflational universe[12] offers ample support to the assumption that classical states of the world emerge around big-bang-like situations as inhomogeneous fluctuations. Taken that as granted, one has to explain how those classical states resist to the kind of quantum spreading described by the linear Schrödinger equation.

Self-sustaining classicality in the above sense needs nonlinearity in the form of an attractive self-interaction acting on the wave function. It can be regarded as a kind of auto-focusing of matter waves, eventually caused by a potential whose source is the matter density. The usual two-point interaction construct of ordinary quantum mechanics is not satisfactory, since it preserves the linearity of the equations of motion, with the spreading of wave packets, formation of Schrödinger cats and all that. On the other hand, *mean-field approximation* to a two-point attractive interaction is just the thing we need to keep wave packets together. There is a trouble though: for the usual kind of interactions, mean field is an approximation, not the true nature of things.

In all respects, gravity is the natural candidate[13]. It is none of the familiar class of contact interactions, since it is encoded into the geometry of space-time. There is no evidence against *defining* mass density, the source of gravity, as the quantum mechanical mean value of the operator of mass density, without additional quantum fluctuations. In that case there is no quantum fluctuation of the metric either, and gravity acts as a mean field on all massive objects, including its sources[14]. Such a scheme in the limiting case of Newtonian gravity was described in Diósi (1984)[15], who – following the protocol of Bialynicki-Birula and Mycielski[16] – demonstrated the existence of soliton-like wave packets stabilized by gravitation, pointing out that gravitation is not able to pull distinct wave packets together.

However, that is not necessary for self-sustaining classicality. We suggest a scenario in which although Eq. (1) remains valid, the two apparatus wave functions ψ_+ and ψ_- remain close to each other, glued together by gravitation, although the measurement interaction tends to pull them apart. As a consequence, the double wave packet, slightly split to balance between opposing forces, moves together under the action of the averaged measurement force, weighted by $|c_+|^2$ and $|c_-|^2$ in a mean-field way; then entering a frozen random environment, undergoes a random choice with the respective probabilities p and $1 - p$, in agreement with Born's rule.

As demonstrated below in detail, for the above scenario gravitation has to be strong enough to protect a double wave packet from splitting under the action of a measuring interaction which is strong enough to accelerate it as a whole.

To simplify the analysis, we start from the two-state

microobject interacting with a one-dimensional massive apparatus, introduced in Eq. (1). The model Hamiltonian is

$$\hat{H} = |+\rangle\langle+| \otimes \hat{h}_+ + |-\rangle\langle-| \otimes \hat{h}_-, \quad (2)$$

where

$$\hat{h}_\pm = -\frac{\hbar^2}{2M}\partial_x^2 + V_{grav}(x, t) \mp F_{meas} \cdot x - F_{div} \cdot x \quad (3)$$

contains a constant force $\pm F_{meas}$ representing the interaction of the microobject in state $|\pm\rangle$ with the apparatus, as well as a force F_{div} modelling a landscape of random barriers, frozen during the act of measurement, diverting the apparatus towards one of the possible outcomes of the measurement.

We set out to solve the time-dependent Schrödinger equation for the pure quantum state on the r.h.s. of Eq. (1). Because of $\langle+|-\rangle = 0$, the c.o.m density of the apparatus is a weighted sum of two non-interfering terms:

$$\varrho(x, t) = p|\psi_+(x, t)|^2 + (1 - p)|\psi_-(x, t)|^2. \quad (4)$$

To calculate the gravitational potential $V_{grav}(x, t)$, we assume that the apparatus is a homogeneous massive sphere of mass M and radius R , and - what should be checked afterwards - both the widths x_0 of the c.o.m wave packets and their distance d are kept small:

$$x_0, d \ll R; \quad (5)$$

then[15]

$$\begin{aligned} V_{grav}(x, t) &= \frac{M\omega_{grav}^2}{2} \int dy (x - y)^2 \varrho(y, t) \\ &= \frac{M\omega_{grav}^2}{2} x^2 - M\omega_{grav}^2 \bar{x}x + \frac{M\omega_{grav}^2}{2} \bar{x}^2. \end{aligned} \quad (6)$$

Here we have introduced the characteristic frequency of an extended object oscillating in its own parabolic gravitational potential,

$$\omega_{grav}^2 = G \frac{M}{R^3}, \quad (7)$$

where G is Newton's gravitational constant. For a typical condensed-matter density $10^4 \text{ Kg } m^{-3}$, one obtains $\omega_{grav} \approx 10^{-3} \text{ Hz}$. We notice that Eq. (6) furnishes a dimensionally correct potential energy that can be plugged into a one-dimensional Schrödinger equation to obtain valid estimates.

The basic estimate is this: to sustain classicality of the apparatus, the double wave packet should remain together during measurement, therefore its mean splitting

should obey $d \approx F_{meas}/M\omega_{grav}^2 < R$. On the other hand, in order to obtain measurement, its displacement as a whole during the measurement interaction of duration τ_{meas} should be large enough: $(F_{meas}/M)\tau_{meas}^2 \geq l_0$, where l_0 is a displacement on the scale of the diverting potential landscape. The combination of the two requirements gives

$$(\omega_{grav}\tau_{meas})^2 > \frac{l_0}{R}. \quad (8)$$

In view of the estimate of ω_{grav} given above, this inequality means that for a measurement of duration 1 s, random decision is controlled on the displacement scale of 10^{-6} times the size of the apparatus. If that scale is fixed about 1 nm, our estimates require $R \gtrsim 10^{-3}$ m for an object big enough to survive conflicting accelerations without splitting into two parts entangled with different states of a microobject. That is our tentative criterion for classicality of the apparatus. Although a massive sphere dragged around by interaction with a microobject is certainly not the realistic model for a particle detector[17], the above figures give sufficient comfort to proceed with the analysis.

Equations (2, 3, 4, 6) define a Hamiltonian depending on the modulus of the wavefunction. However, as seen from Eq. (6), that dependence is only through two time-dependent parameters $\bar{x}^\alpha(t) = \int x^\alpha \varrho(x, t) dx$ ($\alpha = 1, 2$), therefore the corresponding Schrödinger equation, although nonlinear, can be solved by elementary tools.

The tool we use is the equation of motion of a coherent state under the action of a time-dependent force[18], having in mind the time-dependence of the gravitational force proportional to $\bar{x}$ in Eq. (6). The other forces are taken constant in time[19], so we look for a uniformly accelerating motion with $\bar{x} = A + Bt + Ct^2$. Assuming that $\psi_\pm = |\alpha_\pm\rangle$ are coherent states of width $x_0 = (\hbar/M\omega_{grav})^{1/2}$ and respective positions $\bar{x}_\pm = \sqrt{2} x_0 \Re \alpha_\pm$, from which averaging gives

$$\bar{x} = p\bar{x}_+ + (1-p)\bar{x}_- \quad (9)$$

for the moving c.o.m. coordinate. Applying properly scaled formulas from Reference 18, a closed integral equation is obtained for $\bar{x}(t)$. It turns out that as expected, the equation has a smooth, uniformly accelerating solution, the coefficients A, B, C of which can be evaluated, and one obtains

$$\bar{x}(t) = \bar{x}(0) + \bar{v}(0) t + \frac{\mathcal{F}}{2M} t^2, \quad (10)$$

where

$$\mathcal{F} = 2 \left(p - \frac{1}{2}\right) F_{meas} + F_{div} \quad (11)$$

is the sum of the “meanfieldized” measurement force caused by interaction with the microobject, and the frozen random force diverting the motion of the apparatus towards one of the possible outcomes, according to $\mathcal{F} \gtrless 0$. If F_{div} is uniformly distributed between $-F_{meas}$ and F_{meas} , the respective probabilities of moving to the right or to the left are p and $1-p$, in accordance with Born’s rule.

Eq. (10) is a solution appropriate to a special class of initial conditions. For a general initial condition, oscillations around the smooth path appear. We conjecture that the role of decoherence in quantum measurement is to continuously damp such oscillations. Preliminary numerical results seem to support that expectation; details are currently being studied.

What has been described above is a gravity-based dynamical model for a single detector of a binary observable of a microobject. It shows a route to the dynamical origin of quantum randomness, with no creation of macroscopically split superpositions, and accordingly, with no necessity of collapse. Once classical states of matter are present, gravitation forces macroscopic bodies to remain confined, and allows them to explore the possibilities classicality offers to generate randomness. In that respect this model performs like stochastic reduction models[9] do, without putting in randomness by hand.

The present scheme definitely gets out of the Copenhagen philosophy: Eq. (4) materializes the Schrödinger wave function as the source of gravity, which precludes the possibility of relegating the wave function into the realm of ideas.

In one important respect, however, the present approach is less efficient than usual quantum mechanics. This is the rationalization of detector anti-correlations: the feature that for a single particle interacting with two detectors firing on mutually exclusive values of an observable, only one of the detectors can give a signal. That feature, in the root of Heisenberg’s original collapse idea[2], and also very seriously discussed by Einstein[20], is generally believed to hold, although direct experimental tests are not abundant.

In its Neumannian way, ordinary quantum mechanics offers an explanation[21]. In a trivial extension to a two-detector situation in which $\psi_\pm^{(1)}$ means that detector 1 gives a signal or not, and $\psi_\pm^{(2)}$ is the same for detector 2, Eq. (1) is modified like this:

$$\begin{aligned} |\Psi\rangle &= (c_+|+\rangle + c_-|-\rangle)\psi^{(1)}_+(x, 0)\psi^{(2)}_-(x, 0) \\ &\Rightarrow c_+|+\rangle\psi^{(1)}_+(x, t)\psi^{(2)}_-(x, t) \\ &\quad + c_-|-\rangle\psi^{(1)}_-(x, t)\psi^{(2)}_+(x, t). \end{aligned} \quad (12)$$

This equation contains only terms of the expected anti-correlation property, therefore after collapsing into one of the terms of the entangled superposition, a classically ob-

servable anti-correlation would appear between the two detectors.

That nice feature is lost in the present scheme, since $\psi_+^i(x, t)$ and $\psi_-^i(x, t)$ (for $i = 1, 2$) are macroscopically indistinguishable. At the face of our model, the respective probabilities of the two-detector outcomes $++$, $+-$, $-+$, $--$ would be $p(1-p)$, p^2 , $(1-p)^2$, $(1-p)p$. As we all believe to know, the correct answer is $0, p, 1-p, 0$.

That very strongly resembles of interference: destructive on some of the classically apparent possibilities, constructive on others. The trouble is that I was unable to identify the corresponding scenario of interference. One of the less obvious possibilities would be a Berry phase[22] inducing a vector potential in the two-detector configuration space, that would destabilize the $++$ to $--$ diagonal. As a matter of fact, the last term of Eq. (6) gives rise to a fancy phase factor; however, the configuration-space curl of the resulting vector potential is vanishing: no destabilizing force arises. Therefore it is left to the Reader as an exercise to pin down what is interfering with what.

Among the many colleagues with whom I had the pleasure to discuss the issues exposed in this paper, I would like to mention those who left the most significant traces on my thinking: Lajos Diósi and Anton Zeilinger. This work has been partially supported by the Hungarian Research Foundation (grant OTKA T 029544).

-
- [1] M. Born, Z. f. Physik **37**, 863 (1926).
 - [2] W. Heisenberg, *The Physical Principles of the Quantum Theory*, (University of Chicago Press, Chicago, 1930), reprinted by Dover Publications, p. 39.
 - [3] J. von Neumann, *Mathematische Grundlagen der Quantenmechanik*, (Springer, Berlin, 1932); English translation: *Mathematical Foundations of Quantum Mechanics*, (Princeton University Press, 1955).
 - [4] The acknowledged classics on related issues is *Quantum Theory and Measurement*, J. A. Wheeler and W. H. Zurek, eds. (Princeton University Press, Princeton 1983).
 - [5] J. S. Bell, Physics World **3**, 33 (1990).
 - [6] H. D. Zeh, Found. Phys. **1**, 69 (1970); W. H. Zurek, Phys. Rev. D **24**, 1516 (1981).
 - [7] For a recent extensive review, see D. Giulini, E. Joos, C. Kiefer, J. Kupsch, I.-O. Stamatescu, H. D. Zeh, *Decoherence and the Appearance of a Classical World in Quantum Theory*, (Springer, Berlin, 1996).
 - [8] This is an issue of quantum computing; see e. g. D. Bacon, J. Kempe, D. A. Lidar and K. B. Whaley, Phys. Rev. Lett. **85**, 1758 (2000); A. Beige, D. Braun and P. L. Knight, Phys. Rev. Lett. **85**, 1762 (2000).
 - [9] G. C. Ghirardi, A. Rimini and T. Weber, Il Nuovo Cimento **30B**, 133 (1975); P. Pearle, Phys. Rev. A **39**, 2277 (1989); L. Diósi, Phys. Lett. A **129**, 419 (1988); see also Ref.7.
 - [10] Y. Aharonov, D. Z. Albert and L. Vaidman, Phys. Rev. Lett. **60**, 1351 (1988).
 - [11] Using a conceptual framework introduced by A. Peres, Phys. Rev. A **30**, 1610 (1984), the effect was first described by R. Jalabert and H. M. Pastawski, Phys. Rev. Lett. **86**, 2490 (2001), and further analyzed by Ph. Jacquod, P. G. Silvestrov and C. W. J. Beenakker, Phys. Rev. E **64**, 055203(R) (2001).
 - [12] For a readable account, see e. g. A. D. Linde, *Inflation and Quantum Cosmology*, (Academic Press, Boston, 1990).
 - [13] Connecting gravitation with wave-vector collapse has a long history. Among the significant contributions, we mention R. P. Feynman, *Lectures on Gravitation*, (California Institute of Technology, Berkeley, 1962-1963); F. Károlyházi, Nuovo Cimento A **42**, 390 (1966); F. Károlyházi, A. Frenkel and B. Lukács, in: *Quantum Concepts in Space and Time*, R. Penrose and C. J. Isham, eds. (Oxford University Press, Oxford, 1986) p. 109; L. Diósi, Phys. Rev. A **40**, 1165 (1989); G. C. Ghirardi, R. Grassi and A. Rimini, Phys. Rev. A **42**, 1057 (1990); R. Penrose, General Relativity and Gravitation **28**, 581 (1996); J. Anandan, Found. Phys. **29**, 333-348 (1999).
 - [14] Evidence against the scheme would be detecting gravity-induced quantum correlations, like the ones of coulombic origin, familiar in quantum chemistry; obviously an experimental check is by many orders of magnitude beyond feasibility.
 - [15] L. Diósi, Phys. Lett. A **105**, 199 (1984).
 - [16] I. Białynicki-Birula and J. Mycielski, Ann. Phys. (N.Y.) **100**, 62 (1976).
 - [17] In any of the realistic cases, indirect gravitational confinement is expected to dominate, e.g. electrons of a photodetector confined in the coulombic cage of a gravitationally confined crystal.
 - [18] See e.g. C. W. Gardiner: *Quantum Noise*, (Springer, Berlin, 1991) pp. 115-116.
 - [19] The assumption of constant external forces is not essential: a short-time expansion for forces and c.o.m. is converted into a high-frequency asymptotic series for $\alpha_{\pm}(t)$, whose coefficients can be calculated in succession.
 - [20] A. Einstein, *Fifth Solvay Conference, 1927*, discussed in: M. Jammer, *The Philosophy of Quantum Mechanics*, (Wiley-Interscience Publications, New York, 1974), pp. 115-117.
 - [21] That explanation seems to have been overlooked by authors of papers on “one-particle nonlocality”: see L. Hardy, Phys. Rev. Lett. **73**, 279 (1994); with related comments by L. Vaidman, and by D. M. Greenberger, M. A. Horne and A. Zeilinger, and reply by L. Hardy: *ibid.* **75**, 2063, 2064, 2065 (1995).
 - [22] M. V. Berry, Proc. Roy. Soc. A **392**, 45 (1984); C. A. Mead, Rev. Mod. Phys. **64**, 51 (1992); Y. Aharonov, T. Kaufherr, S. Popescu and B. Reznik, Phys. Rev. Lett. **80**, 2023 (1998).